

# Anthracnose Fruit Rot of Strawberry<sup>1</sup>

James C. Mertely, Bruna B. Forcelini, and Natalia A. Peres<sup>2</sup>

Anthracnose fruit rot, caused by the fungus *Colletotrichum acutatum*, is an important disease for strawberry world-wide. Other species of *Colletotrichum*, such as *C. fragariae* and *C. gloeosporioides*, are less frequently involved in fruit rot. Although fruit are most frequently affected by *C. acutatum*, other organs of the plant, including flowers, crowns, leaves, petioles, and roots, are also susceptible.

## Pathogens and Symptoms

Anthracnose fruit rot lesions appear as dark, sunken spots on infected fruit (Figure 1). On green fruit, anthracnose lesions are small ( $\frac{1}{16}$ – $\frac{1}{8}$  inch across), hard, sunken, and dark brown or black. Lesions on ripening fruit are larger ( $\frac{1}{8}$ – $\frac{1}{2}$  inch), hard, sunken, and tan-to-dark brown. During wet weather, the lesions become covered with sticky, light orange ooze composed of millions of spores (conidia) in a mucilaginous matrix (Figure 2). When conditions are favorable for infection, multiple lesions nearly cover the fruit, and lesions may appear on petioles (Figure 3). Strawberry flowers are highly susceptible and turn brown and remain attached to the plant when infected (Figure 4). Flowers affected by the gray mold fungus *Botrytis cinerea* may show similar symptoms. Small black spots on green button-sized fruit may also develop from flower infections (Figure 5).



Figure 1. Anthracnose lesions on a ripe fruit

Credits: UF/IFAS GCREC

## Disease Development and Spread

The most common way a new strawberry crop is infected in annual production systems such as Florida is through transplants from the nursery. *C. acutatum* is a strong colonizer of runner plants in the nursery and may be

1. This document is PP-207, one of a series of the Plant Pathology Department, UF/IFAS Extension. Original publication date March 2005. Revised January 2009, September 2012, and August 2017. Visit the EDIS website at <http://edis.ifas.ufl.edu>.
2. James C. Mertely, research coordinator; Bruna B. Forcelini, graduate assistant; and Natalia A. Peres, professor, Plant Pathology Department, UF/IFAS Gulf Coast Research and Education Center, Wimauma, FL 33598.

The use of trade names in this publication is solely for the purpose of providing specific information. UF/IFAS does not guarantee or warranty the products named, and references to them in this publication do not signify our approval to the exclusion of other products of suitable composition.

The Institute of Food and Agricultural Sciences (IFAS) is an Equal Opportunity Institution authorized to provide research, educational information and other services only to individuals and institutions that function with non-discrimination with respect to race, creed, color, religion, age, disability, sex, sexual orientation, marital status, national origin, political opinions or affiliations. For more information on obtaining other UF/IFAS Extension publications, contact your county's UF/IFAS Extension office.

U.S. Department of Agriculture, UF/IFAS Extension Service, University of Florida, IFAS, Florida A & M University Cooperative Extension Program, and Boards of County Commissioners Cooperating. Nick T. Place, dean for UF/IFAS Extension.

present on the foliage of apparently healthy plants. In some cases, infected runner plants may show visible lesions on the petioles (Figure 3) and roots. A new crop may also be infected by inoculum carried over from the previous crop. In the past, traditional post-season crop destruction and cultivation were believed to eliminate inoculum carryover from Florida production fields. However, *C. acutatum* has been recovered from dead plants left on old plastic during the summer. Thus, if strawberry is planted on old plastic, the inoculum from the old plants could affect the new crop. Various weeds in and around production fields may also be colonized by *C. acutatum* from strawberry.



Figure 2. Spore mass of *C. acutatum* on anthracnose lesion  
Credits: UF/IFAS GCREC



Figure 4. Anthracnose flower blight  
Credits: UF/IFAS GCREC




Figure 5. Anthracnose lesion on green fruit  
Credits: UF/IFAS GCREC



Figure 3. Anthracnose lesions on petioles  
Credits: UF/IFAS GCREC

*C. acutatum* appears to spread first on the foliage, often without causing visible symptoms. A few conidia (asexual spores) are formed on green leaves and petioles, and more are produced as the tissue ages and dies. Conidia are moved from the foliage to the flowers and fruit primarily by splashing water. They then germinate and infect. Developing infections on flowers and fruit produce abundant conidia that are spread to other plants and fields by equipment and harvesters, especially when the plants are wet.

Anthracnose fruit rot development is favored by warm, wet weather. For this reason, epidemics typically occur in the spring when conditions are more favorable. Such epidemics



cause serious losses for Florida growers. Crop losses occur mostly in the field since visibly spotted fruit are culled and the development of latent infections postharvest is suppressed by precooling and refrigeration.

## Control

Anthracnose fruit rot is best controlled by exclusion (i.e., by not introducing the pathogen into the field). Wherever possible, transplants should be obtained from pathogen-free nurseries. In addition, moving personnel and equipment from diseased to healthy fields without proper cleaning should be avoided. All cultivars recently released in Florida ('Florida 127', 'Winterstar', and 'Florida Radiance') have higher levels of resistance to anthracnose than 'Strawberry Festival'. When highly susceptible cultivars are grown, regular preventive fungicide applications are needed to suppress the disease.

In central Florida, anthracnose management has been based on weekly applications of the broad-spectrum, protectant fungicide captan. Applications can be made at low label rates early in the season because weather conditions are less favorable for disease development at that time. Often a few anthracnose-infected flowers and fruit that appear in late January or early February lead to epidemics during the main harvest in late February and March. During the critical January to March period, protectant fungicides should be applied at higher label rates, and additional fungicides with some curative activity, such as the strobilurin fungicides, Abound® and Cabrio®, or Switch®, may be needed for adequate disease control. However, the emergence of *C. acutatum* populations resistant to the strobilurin fungicides (Fungicide group 11) was first reported in Florida in 2013. Since then, lack of AFR control has been observed in commercial fields where resistant populations of *C. acutatum* are present. It is recommended that growers with AFR-symptomatic plants send samples to the UF-GCREC plant disease clinic to determine whether plants harbor resistant *C. acutatum*. If resistance is detected, AFR control should rely mainly on preventive applications of captan.

Since anthracnose incidence is highly dependent on weather conditions, fungicide applications can be timed based on key factors, such as leaf wetness duration and temperature during the wetting period. The Strawberry Advisory System (StAS) (<http://edis.ifas.ufl.edu/ae450>) (<http://www.agroclimate.org/tools/strawberry/fl>) has been developed to advise growers of the need to spray to prevent anthracnose as well as Botrytis fruit rot. StAS provides growers with recommendations for timing as well as

suggestions for selection of the most appropriate product. Disease risk alerts can also be provided to growers via e-mail or text messages.

Research has demonstrated that fungicide sprays can be reduced without affecting disease control or yield when following the StAS recommendations, especially when the more tolerant cultivars, such as Florida 127, Florida Radiance, and Winterstar are grown.

A list of fungicide products recommended for anthracnose fruit rot control based on research trials is provided in Table 1.

Table 1. Fungicides recommended for control of Anthracnose fruit rot of strawberries in Florida

Product name (active ingredient)	Fungicide group <sup>a</sup>	Maximum rate/A		Minimum days to harvest	Re-entry interval (hours)	Remarks
		Per application	Per season			
Captan 80 WDG (captan)	M4	3.75 lb.	30 lb.	1	24	Do not apply in combination with or immediately before or closely following oil sprays. Do not mix with strongly alkaline materials.
Captec 4L® (captan)	M4	3 qt.	54 qt.	1	24	Do not apply in combination with or immediately before or closely following oil sprays. Do not mix with strongly alkaline materials.
Quilt Xcel® (propiconazole + azoxystrobin) <sup>b</sup>	3 + 11	14 fl. oz.	56 fl. oz.	0	12	Do not make more than two consecutive applications before alternating to another fungicide group and no more than four applications per crop per year.
Merivon (fluxapyroxad + pyraclostrobin) <sup>b</sup>	7 + 11	11 fl oz.	33 fl oz.	0	12	Do not make more than two sequential applications before alternating to another non-group 7 or non-group 11 fungicide
Abound® (azoxystrobin) <sup>b</sup>	11	15.4 fl. oz.	1.92 qt.	0	4	Do not make more than two consecutive applications and no more than four applications per crop per year.
Cabrio® EG (pyraclostrobin) <sup>b</sup>	11	14 fl. oz.	70 fl. oz.	0	12	Do not make more than two consecutive applications and no more than five applications per crop per year.
Evito® 480 SC, Aftershock™ (fluoxastrobin) <sup>b</sup>	11	5.7 fl. oz.	22.8 fl. oz.	1	12	Do not make more than two consecutive applications and no more than four applications per crop per year.
Flint® (trifloxystrobin) <sup>b</sup>	11	3.2 oz.	19.2 oz.	1	12	Do not make more than two consecutive applications and no more than six applications of group 11 fungicides per crop per year.
Switch® 62.5 WG (cyprodinil + fludioxonil)	9 + 12	14 oz.	56 oz.	0	12	Do not make more than two consecutive applications. Do not plant crops not on the label for 30 days after last application.

<sup>a</sup> Fungicide group (FRAC Code): Numbers (1–37) and letters (M) are used to distinguish the fungicidal mode of action groups. All fungicides within the same group (with same number or letter) indicate same active ingredient or similar mode of action. This information must be considered when making decisions about how to manage fungicide resistance. M = multisite inhibitors, fungicide resistance is low; NC = not classified. Source: <http://www.frac.info/> (Fungicide Resistance Action Committee, FRAC).

<sup>b</sup> Resistance reported to group 11 fungicides in *C. acutatum* isolates from strawberry. Samples can be submitted to the UF/IFAS GCREC strawberry lab to determine if resistance is present in certain fields. **Always read a current product label before applying any chemicals.**